

2015 New York Prevention Needs Assessment Survey



Survey Results for:
Jefferson County

Sponsored by:
Pivot Prevention and Health Services
167 Polk Street, Suite 320
Watertown, NY 13601

2015 Prevention Needs Assessment Survey Profile Report

This report summarizes the findings from the Prevention Needs Assessment (PNA) Survey that was conducted during 2015. The results are presented along with comparisons to national data sources such as the Monitoring the Future Survey (only grades 8, 10, and 12 are surveyed) and the Bach Harrison Norm (BH Norm), which consists of a large, weighted, nationwide sample.

The survey was designed to assess students' involvement in a specific set of problem behaviors, as well as their exposure to a set of scientifically validated risk and protective factors. The risk and protective factors have been shown to influence the likelihood of academic success, school dropout, substance abuse, violence, and delinquency among youth.

Table 1 contains the characteristics of the students who completed the survey from your community. When using the information in this report, please pay attention to the number and

Contents:

Introduction
The Risk and Protective Factor Model of Substance Abuse Prevention
Building a Strategic Prevention Framework
Validity Measures
How to Read the Charts
Tools for Assessment and Planning
ATOD and Antisocial Behavior Charts
Risk and Protective Factor Charts
Risk and Protective Factor Scale Definitions
Data Tables

Drug Free Communities and Youth Perception of Substance Use Report
Contacts for Prevention

The Risk and Protective Factor Model of Substance Abuse Prevention

percentage of students who participated from your community. The sample size for this survey administration was 6,803 students. If 60% or more of the students sample participated, the report is a good indicator of the levels of substance use, risk, protection, and antisocial behavior. If fewer than 60% participated, a review of who participated should be completed prior to generalizing the results to the entire community.

Many states and local agencies have adopted the Risk and Protective Factor Model to guide their prevention efforts. The Risk and Protective Factor Model of Prevention is based on the simple premise that to prevent a problem from happening, we need to identify the factors that increase the risk of that problem developing and then find ways to reduce the risks. Just as medical researchers have found risk factors for heart disease such as diets high in fat, lack of exercise, and smoking; a team of researchers at the University of Washington have defined a set of risk factors for youth problem behaviors.

Risk factors are characteristics of school, community, and family environments, as well as characteristics of students and their peer groups that are known to predict increased likelihood of drug use, delinquency, school dropout, teen pregnancy, and violent behavior among youth. Dr. J. David Hawkins, Dr. Richard F. Catalano, and their colleagues at the Development Research Group have investigated the relationship between risk and protective factors and youth problem behavior. For example, they

Table 1. Characteristics of Participants

Table 1. Characteristics of Participants					
Student Totals					
Jefferson County					
Total Students	2012		2014		2015
	Number	Percent	Number	Percent	Number
				Percent	
	2735	100	2103	100	5571
Grade					
7	n/a	n/a	n/a	n/a	963
8	1056	38.6	861	40.9	890
9	n/a	n/a	n/a	n/a	994
10	964	35.2	724	34.4	920
11	n/a	n/a	n/a	n/a	860
12	715	26.1	518	24.6	944
Gender					
Male	1324	48.7	1052	50.4	2761
Female	1397	51.3	1037	49.6	2777
Ethnicity					
Native American	45	1.7	32	1.5	97
Asian	35	1.3	36	1.7	98
African American	168	6.2	133	6.4	329
Pacific Islander	12	0.4	15	0.7	48
Hispanic	125	4.6	152	7.3	4082
White	2074	76.2	1525	73.3	603
Multi-racial or Other	263	9.7	217	10.4	740
					10.9

Table 1. represents the total survey population. Students were given the option to skip questions, and not all students completed the survey. The percentages in remaining tables/figures of this report reflect the percent of students responding to each question, rather than the percent of the total survey population.

The Risk and Protective Factor Model of Substance Abuse Prevention (Continued)

have found that children who live in families with high levels of conflict are more likely to become involved in problem behaviors such as delinquency and drug use than children who live in families with low levels of family conflict.

Protective factors exert a positive influence or buffer against the negative influence of risk, thus reducing the likelihood that adolescents will engage in problem behaviors. Protective factors identified through research reviewed by Drs. Hawkins and Catalano include social bonding to family, school, community and peers; healthy beliefs and clear standards for behavior; and individual characteristics. For bonding to serve as a protective influence, it must occur through involvement with peers and adults who communicate healthy values and set clear standards for behavior. Research on risk and protective factors has important implications for prevention efforts.

The premise of this approach is that in order to promote positive youth development and prevent problem behaviors, it is necessary to address those factors that predict the problem.

By measuring risk and protective factors in a population, prevention programs can be implemented that will reduce the elevated risk factors and increase the protective factors. For example, if academic failure is identified as an elevated risk factor in a community, then mentoring, tutoring, and increased opportunities and rewards for classroom participation can be provided to improve academic performance. The chart to the right shows the links between the 20 risk factors and the six problem behaviors placed in the chart to indicate where at least two well designed, published research studies have shown a link between the risk factor and the problem behavior.

Risk Factors for Adolescent Problem Behavior						Community					
Problem Behaviors	Substance Abuse	Delinquency	Teen Pregnancy	School Drop-Out	Violence	Depression & Anxiety	Availability of Drugs	Availability of Firearms	Community Laws and Norms	Favorable Toward Drug Use, Firearms and Crime	Media Portrayals of the Behavior
	✓						✓				✓
								✓			
Family											
	✓	✓	✓	✓	✓	✓	Family History of the Problem Behavior	✓	✓	✓	✓
	✓	✓	✓	✓	✓	✓	Family Management Problems	✓	✓	✓	✓
	✓	✓	✓	✓	✓	✓	Family Conflict	✓	✓	✓	✓
	✓	✓	✓	✓	✓	✓	Favorable Parental Attitudes and Involvement in the Problem Behavior	✓	✓	✓	✓
School											
	✓	✓	✓	✓	✓	✓	Academic Failure Beginning in Late Elementary School	✓	✓	✓	✓
	✓	✓	✓	✓	✓	✓	Lack of Commitment to School	✓	✓	✓	✓
Peer / Individual											
	✓	✓	✓	✓	✓	✓	Early & Persistent Antisocial Behavior	✓	✓	✓	✓
	✓	✓	✓	✓	✓	✓	Rebelliousness	✓	✓	✓	✓
	✓	✓	✓	✓	✓	✓	Gang Involvement	✓	✓	✓	✓
	✓	✓	✓	✓	✓	✓	Friends Who Engage in the Problem Behavior	✓	✓	✓	✓
	✓	✓	✓	✓	✓	✓	Favorable Attitudes Toward the Problem Behavior	✓	✓	✓	✓
	✓	✓	✓	✓	✓	✓	Early Imitation of the Problem Behavior	✓	✓	✓	✓
	✓	✓	✓	✓	✓	✓	Constitutional Factors	✓	✓	✓	✓

The check marks have been placed in the chart to indicate where at least two well designed, published research studies have shown a link between the risk factor and the problem behavior.

Building a Strategic Prevention Framework

The survey is an important data source for the Substance Abuse and Mental Health Services Administration (SAMHSA) Center for Substance Abuse Prevention (CSAP) Strategic Prevention Framework (SPF). CSAP created the SPF model to guide states and communities in creating planned, data-driven, effective, and sustainable prevention programs. Each part represents an interdependent element of the ongoing process of prevention coordination.


Assessment: Profile Population Needs, Resources, and Readiness to Address the Problems and Gaps in Service Delivery. The SPF begins with an assessment of the needs in the community that is based on data. One of the primary sources of needs assessment data is this Prevention Needs Assessment Survey (PNA). While planning prevention services, communities are urged to collect and use multiple data sources, including archival and social indicators, assessment of existing resources, key informant interviews, and community readiness. The PNA results presented in this Profile Report will help you to identify needs for prevention services. PNA data include adolescent substance use, anti-social behavior, and many of the risk and protective factors that predict adolescent problem behaviors.

Capacity: Mobilize and/or Build Capacity to Address Needs. Engagement of key stakeholders at the State and community levels is critical to plan and implement successful prevention activities that will be sustained over time. Some of the key tasks to mobilize the state and communities are to work with leaders and stakeholders to build coalitions, provide training, leverage resources, and help sustain prevention activities.

Planning: Develop a Comprehensive Strategic Plan

States and communities should develop a strategic plan that articulates not only a vision for the prevention activities, but also strategies for organizing and implementing prevention efforts. The strategic plan should be based on the assessments conducted during Step 1. The Plan should address the priority needs, build on identified resources/strengths, set measurable objectives, and identify how progress will be monitored. Plans should be adjusted with ongoing needs assessment and monitoring activities.

Implementation: Implement Evidence-based Prevention Programs and Infrastructure Development Activities. By measuring and identifying the risk factors and other causal factors that contribute to the targeted problems specified in your strategic plan, programs can be implemented that will reduce the prioritized substance abuse problems. After completing Steps 1, 2, and 3, communities will be able to choose prevention strategies that have been shown to be effective, are appropriate for the population served, can be implemented with fidelity, are



Building a Strategic Prevention Framework (cont'd)

culturally appropriate, and can be sustained over time. The Western Center for the Application of Prevention Technology has developed an internet tool located at <http://casat.unr.edu/bspractices/research.php> for identifying Best Practice Programs. Another resource for evidence-based prevention practices is SAMHSA's National Registry of Evidence-based Programs and Practices www.nrcpp.samhsa.gov.

Evaluation: Monitor Process, Evaluate Effectiveness, Sustain Effective Programs/Activities, and Improve or Replace Those That Fail. Finally, ongoing monitoring and evaluation are essential to determine if the desired outcomes are achieved, assess service delivery quality, identify successes, encourage needed improvement, and promote sustainability of effective policies, programs, and practices. The OPNA allows communities to monitor levels of ATOD use, antisocial behavior, risk, and protection.

Sustainability and Cultural Competence: Incorporate principles of cultural competence and sustainability in each of the five elements. At the center of the SPF model, sustainability and cultural competence play a key role in assessment, capacity appraisal, planning, implementation and evaluation, ensuring successful, long lasting prevention programs.

Sustainability is accomplished by utilizing a comprehensive approach. States and communities should plan adaptive, flexible programs around a variety of resources, funding, and organizations. An inclusive design helps build sustainable programs and achieve sustainable outcomes. A strategic plan that dynamically responds to changing issues, data, priorities, and resources is more likely to achieve long term results.

Sharing information gathered during the evaluation stage with key stakeholders, forging partnerships and encouraging creative collaboration all enhance sustainability.

Cultural Competence recognizes unique needs, styles, values and beliefs of the recipients of prevention efforts. Culturally competent prevention strategies use interventions, evaluations and communication strategies appropriate to their intended community. Cultural issues reflect a range of influences and are not just a matter of ethnic or racial identity. Learning to communicate with audiences from diverse geographic, cultural, economic, social, and linguistic backgrounds can increase program efficacy and ensure sustainable results.

Whether enlisting extended family networks as a prevention resource for single parent households, or ensuring there are resources available to bridge language gaps, cultural competency will help you recognize differences in prevention needs and tailor prevention approaches accordingly.

A one-size-fits-all program is less effective than a program that draws on community-based values, traditions, and customs and works with knowledgeable people from the community to develop focused interventions, communication, and support.

Validity Measures

Honesty: Because the survey was anonymous, and because confidentiality was stressed through the survey's administration process, most of the reasons for students to exaggerate or deny behaviors were eliminated. However, Bach Harrison has built several checks into the data analysis to minimize the impact of students who were either not truthful in their responses or who did not take the survey seriously. Surveys were eliminated from the final data reported in this report for meeting one or more the following five pre-determined dishonesty indicators:

1. In response to a question about whether or not they had been honest in completing the survey, the students indicated that they were "Not Honest At All" in completing the survey.
2. The students indicated that they had used a non-existent, fictitious drug in their lifetime or in the past 30 days.
3. The students reported an impossibly high level of multiple drug use (having used substances on 120 or more occasions in the past 30 days).
4. The students indicated past-month use rates that were higher than lifetime use rates.

5. The students reported an age that was inconsistent with their grade or their school; for example, a 10 year-old 12th grader or 19 year old 6th grader.
- Additionally, if a student did not answer enough of the validity questions to determine whether or not they were honest in their responses, their survey data were also removed from the final analysis presented in this report.

There are four types of charts presented in this report:

1. Substance use charts
2. Antisocial behavior (ASB) and Gambling charts
3. Risk factor charts
4. Protective factor charts.

Data from the charts are also presented in Tables 3 through 10. Additional data found in later tables are explained at the end of this section.

Understanding the Format of the Charts

There are several graphical elements common to all the charts. Understanding the format of the charts and what these elements represent is essential in interpreting the results of the PNA survey.

The Bars on substance use and antisocial behavior charts represent the percentage of students in that grade who reported a given behavior. The bars on the risk and protective factor charts represent the percentage of students whose answers reflect significant risk or protection in that category. Each set of differently colored bars represents one of the past administrations of the PNA. By looking at the percentages over time, it is possible to identify trends in substance use and antisocial behavior. By studying the percentage of youth at risk and with protection over time, it is possible to determine whether the percentage of students at risk or with protection is increasing, decreasing, or staying the same. This information is important when deciding which risk and protective factors warrant attention.

Dots and Diamonds provide points of comparison to larger samples. The dots on the charts represent the percentage of all of the youth surveyed who reported substance use, problem behavior, elevated risk, or elevated protection. Please note that the dot represents the aggregate results of all participating students rather than a random sample of students. The survey results provide considerable information for communities to use in planning prevention services.

The diamonds represent national data from either the Monitoring the Future (MTF) Survey or the Bach Harrison Norm (BH Norm). The BH Norm was developed by Bach Harrison L.L.C. to provide states and communities with the ability to compare their results on risk, protection, and antisocial measures with more national measures. Survey participants from eight statewide surveys and five large regional surveys across the nation were combined into a database of approximately 460,000 students. The results were weighted to make the contribution of each state and region proportional to its share of the national population. Bach Harrison analysts then calculated rates for antisocial behavior and for students at risk and with protection. The results appear on the charts as BH Norm. In order to keep the BH Norm relevant, it is updated approximately every two years as new data become available.

A comparison to state-wide and national results provides additional information for your community in determining the relative importance of levels of alcohol, tobacco and other drug (ATOD) use, antisocial behavior, risk, and protection. Information about other students in the state and the nation can be helpful in determining the seriousness of a given level of problem behavior. Scanning across the charts, it is important to observe the factors that differ the most from the BH Norm. This is the first step in identifying the levels of risk and protection that are higher or lower than those in other communities. The risk factors that are higher than the BH Norm and the protective factors are lower than the BH Norm are probably the factors that you should consider addressing when planning prevention programs.

Cut-Points

Before the percentage of youth at risk on a given scale could be calculated, a scale value or cut-point needed to be determined that would separate the at-risk group from the not at-risk group. The Prevention Needs Assessment (PNA) survey was designed to assess adolescent substance use, anti-social behavior, and the risk and protective factors that predict these adolescent problem behaviors. Since the PNA survey has recently been given to over 460,000 youth nationwide, it was possible to select two groups of youth, one that was more at risk for problem behaviors and another group that was less at risk. A cut-point score was then determined for each risk and protective factor scale that best divided the youth from the two groups into their

Antisocial Behavior, Driving and Alcohol, and Gambling Charts

Antisocial behavior (ASB) is a measure of the percentage of students who report any involvement during the past year with the two antisocial behaviors listed in the charts.

Driving and Alcohol is a measure of the percentage of students who report drinking and driving, or being a passenger in a car where the driver had been drinking in the past 30 days.

Gambling Behavior is a measure of the percentage of students who report any involvement during the past year with the ten types of gambling listed in the charts. *Gambled in the Past Year* is a measure of any participation in any of the gambling types whatsoever.

Risk and Protective Factor Charts

Risk and protective factor scales measure specific aspects of a youth's life experience that predict whether he/she will engage in problem behaviors. The scales, defined in Table 2, are grouped into four domains: community, family, school, and peer/individual. The risk and protective factor charts show the percentage of students at risk and with protection for each of the scales. Along with the scales, there are bars that show the percentage of High Risk Youth and percentage of High Protection Youth. High Risk Youth is defined as the percentage of students who have more than a specified number of risk factors operating in their lives. The number of factors is listed on the charts and tables. High Protection Youth is defined as the percentage of students who have more than a specified number of protective factors operating in their lives and is also listed on the tables and graphs.

The appropriate group, more at-risk or less at-risk. The criteria for separating youth into the more at-risk and the less at-risk groups included academic grades (the more at-risk group received "D" and "F" grades, the less at-risk group received "A" and "B" grades), ATOD use (the more at-risk group had more regular use, the less at-risk group had no drug use and use of alcohol or tobacco on only a few occasions), and antisocial behavior (the more at-risk group had two or more serious delinquent acts in the past year, the less at-risk group had no serious delinquent acts).

The cut-points that were determined by analyzing the results of the more at-risk and less at-risk groups will remain constant and will be used to produce the profiles for future surveys.

Since the cut-points for each scale will remain fixed, the percentage of youth above the cut-point on a scale (at-risk) will provide a method for evaluating the progress of prevention programs over time. For example, if the percentage of youth at risk for family conflict in a community prior to implementing a community-wide family/parenting program was 60% and then decreased to 50% one year after the program was implemented, the program would be viewed as helping to reduce family conflict.

Lifetime, 30 Day & Heavy ATOD Use Charts

There are three types of use measured on the ATOD charts.

Ever-used is a measure of the percentage of students who used the particular substance at least once in their lifetime and is used to show the percentage of students who have had experience with a particular substance.

30-day use is a measure of the percentage of students who used the substance at least once in the 30 days prior to taking the survey and is a more sensitive indicator of the level of current use of the substance.

Heavy use is measured in two ways: *binge drinking* (five or more drinks in a row over the last two weeks), and use of *one-half a pack or more of cigarettes per day*.

Additional Tables in this Report

Table 11 presents the percentages of how and where students obtained and used alcohol during the past year. The data focus on a subgroup of students who indicated at least one means of obtaining or using alcohol. (Students reporting no alcohol use are not represented.) It is important to note that the table represents a subgroup of users and not the entire survey population. Additionally, the smaller the sample, the more dramatic the influence of a student's responses. For example, if only one student in a particular grade reported where he/she obtained alcohol, each category would show up as either 0% or 100%. The table indicates the sample size for each grade surveyed to help clarify the value of the data.

After the Student Alcohol Tables are tables containing information required by communities with CSAP Grants, such as the parent attitudes regarding drinking, police response to drinking, and problems associated with drinking.

After the CSAP questions are tables containing information required by communities with Drug Free Communities Grants, such as the perception of the risks of ATOD use, perception of parent and peer disapproval of ATOD use, past 30-day use, and average age of first use.

After the DFC Tables are the Youth Perception Tables. Youth often overestimate the percentage of their peers who are using substances. Youth perceptions of the percentage of their peers who use cigarettes, alcohol, marijuana, and other illegal drugs are shown in these tables.

Finally, there are any extra questions your agency might have asked.

No Child Left Behind

1. The Safe and Drug Free Schools and Communities section of the No Child Left Behind Act (NCLB) requires that schools and communities use guidelines in choosing and implementing federally funded prevention and intervention programs. The results of the PNA Survey presented in this report can help your schools and community comply with the NCLB Act in three ways:
 1. Programs must be chosen based on objective data about problem behaviors in the communities served. The PNA reports these data in the substance use and antisocial behavior charts and tables presented on the following pages.
2. NCLB-approved prevention programs can address not only substance use and antisocial behavior (ASB) outcomes, but also behaviors and attitudes demonstrated to be predictive of the youth problem behaviors. Risk and protective factor data from this report provide valuable information for choosing prevention programs.
3. Periodic evaluations of outcome measures must be conducted to evaluate the efficacy of ongoing programs. This report provides schools and communities the ability to compare past and present substance use and ASB data.

What are the numbers telling you?

Review the charts and data tables presented in this report. Note your findings as you discuss the following questions.

Which 3-5 risk factors appear to be higher than you would want when compared to the Bach Harrison Norm?

Which 3-5 protective factors appear to be lower than you would want when compared to the Bach Harrison Norm?

Which levels of 30-day drug use are increasing and/or unacceptably high? Which substances are your students using the most? At which grades do you see unacceptable usage levels?

Which antisocial behaviors are increasing and/or unacceptably high? Which behaviors are your students exhibiting the most? At which grades do you see unacceptable behavior levels?

How to identify high priority problem areas
Once you have familiarized yourself with the data, you can begin to identify priorities.

Look across the charts for items that stand out as either much higher or much lower than the others. Compare your data with statewide, and/or national data. Differences of 5% between local and other data are probably significant.

Prioritize problems for your area according to the issues you've identified. Which can be realistically addressed with the funding available to your community? Which problems fit best with the prevention resources at hand?

Determine the standards and values held within your community. For example: Is it acceptable in your community for a percentage of high school students to drink alcohol regularly as long as that percentage is lower than the overall state rate?

Use these data for planning.

Once priorities are established, use data to guide your prevention efforts.

Substance use and antisocial behavior data are excellent tools to raise awareness about the problems and promote dialogue.

Risk and protective factor data can be used to identify exactly where the community needs to take action.


Promising approaches for any prevention goal are available through resources listed on the last page of this report. These contacts are a great resource for information about programs that have been proven effective in addressing the risk factors that are high in your community, and improving the protective factors that are low.

Sample	Priority Rate 1	Priority Rate 2	Priority Rate 3
6th grad. Fam. Attitude to Drugs (Peer/Indiv. Scale) 0 15% (27 > 8-state av.)			
10th grad. - Rewards for prosocial behavior (School Domain)			
10% down 5% from 2 yrs ago & 16% below state av.)			
8th grad. Burgs Drinking 913% (5% above state av.)			
12th grad. - Drunk/High at School @ 21% (about same as state, but remains a priority)			

Risk Factors
Protective Factors
30-day Substance Abuse
Antisocial Behavior


Substance Use and Antisocial Behavior

LIFETIME, 30 DAY & HEAVY ATOD USE
2015 Jefferson County Student Survey, Grade 7




† Monitoring The Future only surveys grades 8, 10, and 12.

ANTISOCIAL BEHAVIOR AND GAMBLING
2015 Jefferson County Student Survey, Grade 7




Substance Use and Antisocial Behavior

LIFETIME, 30 DAY & HEAVY ATOD USE
2015 Jefferson County Student Survey, Grade 8




** MTF either does not collect data on that substance at all, or not at that level.

ANTISOCIAL BEHAVIOR AND GAMBLING
2015 Jefferson County Student Survey, Grade 8




Substance Use and Antisocial Behavior

LIFETIME, 30 DAY & HEAVY ATOD USE
2015 Jefferson County Student Survey, Grade 9




† Monitoring The Future only surveys grades 8, 10, and 12

ANTISOCIAL BEHAVIOR AND GAMBLING
2015 Jefferson County Student Survey, Grade 9




Substance Use and Antisocial Behavior

2015 Jefferson County Student Survey, Grade 10




** MTF either does not collect data on that substance at all, or not at that level.

2015 Jefferson County Student Survey, Grade 10




Substance Use and Antisocial Behavior

LIFETIME, 30 DAY & HEAVY ATOD USE
2015 Jefferson County Student Survey, Grade 11




† Monitoring The Future only surveys grades 8, 10, and 12.

ANTISOCIAL BEHAVIOR AND GAMBLING
2015 Jefferson County Student Survey, Grade 11




Substance Use and Antisocial Behavior

LIFETIME, 30 DAY & HEAVY ATOD USE
2015 Jefferson County Student Survey, Grade 12




** MTF either does not collect data on that substance at all, or not at that level.

ANTISOCIAL BEHAVIOR AND GAMBLING
2015 Jefferson County Student Survey, Grade 12




Substance Use and Antisocial Behavior

LIFETIME, 30 DAY & HEAVY ATOD USE
2015 Jefferson County Student Survey, All Grades





** MTF either does not collect data on that substance at all, or not at that level.



ANTISOCIAL BEHAVIOR AND GAMBLING
2015 Jefferson County Student Survey, All Grades





Risk and Protective Factor Profiles





Risk and Protective Factor Profiles





Risk and Protective Factor Profiles




Risk and Protective Factor Profiles




Risk and Protective Factor Profiles



Risk and Protective Factor Profiles



High Risk Youth are defined as the percentage of students who have more than a specified number of risk factors operating in their lives: 6th grade: 5 or more risk factors, 7th-9th grades: 6 or more factors, 10th-12th grades: 7 or more factors.



High Protection Youth are defined as youth with: 6th and 7th grades: 3 or more protective factors, 8th-12th grades: 4 or more factors.

Risk and Protective Factor Profiles





Table 2. Scales that Measure the Risk and Protective Factors Shown in the Profiles

<i>Community Domain Risk Factors</i>	
Laws and Norms	Research has shown that legal restrictions on alcohol and tobacco use, such as raising the legal drinking age, restricting smoking in public places, and increased taxation have been followed by decreases in consumption. Moreover, national surveys of high school seniors have shown that shifts in normative attitudes toward drug use have preceded changes in prevalence of use.
Favorable Toward Drug Use	The availability of cigarettes, alcohol, marijuana, and other illegal drugs has been related to the use of these substances by adolescents.
<i>Community Domain Protective Factors</i>	
Rewards for Prosocial Involvement	Rewards for positive participation in activities helps youth bond to the community, thus lowering their risk for substance use.
<i>Family Domain Risk Factors</i>	
Poor Family Management	Parents' use of inconsistent and/or unusually harsh or severe punishment with their children places them at higher risk for substance use and other problem behaviors. Also, parents' failure to provide clear expectations and to monitor their children's behavior makes it more likely that they will engage in drug abuse whether or not there are family drug problems.
Family Conflict	Children raised in families high in conflict, whether or not the child is directly involved in the conflict, appear at risk for both delinquency and drug use.
Sibling Drug Use and Exposure to Adult Antisocial Behavior	When children are raised in a family with a history of problem behaviors (e.g., violence or ATOD use), the children are more likely to engage in these behaviors.
Parental Attitudes Favorable Toward Antisocial Behavior and Parental Attitudes	In families where parents use illegal drugs, are heavy users of alcohol, or are tolerant of children's use, children are more likely to become drug abusers during adolescence. The risk is further increased if parents involve children in their own drug (or alcohol) using behavior, for example, asking the child to light the parent's cigarette or get the parent a beer from the refrigerator.
<i>Family Domain Protective Factors</i>	
Family Attachment	Young people who feel that they are a valued part of their family are less likely to engage in substance use and other problem behaviors.
Opportunities for Prosocial Involvement	Young people who are exposed to more opportunities to participate meaningfully in the responsibilities and activities of the family are less likely to engage in drug use and other problem behaviors.
Rewards for Prosocial Involvement	When parents, siblings, and other family members praise, encourage, and attend to things done well by their child, children are less likely to engage in substance use and problem behaviors.
<i>School Domain Risk Factors</i>	
Academic Failure	Beginning in the late elementary grades (grades 4-6) academic failure increases the risk of both drug abuse and delinquency. It appears that the experience of failure itself, for whatever reasons, increases the risk of problem behaviors.
Low Commitment to School	Surveys of high school seniors have shown that the use of drugs is significantly lower among students who expect to attend college than among those who do not. Factors such as liking school, spending time on homework, and perceiving the coursework as relevant are also negatively related to

Risk and Protective Scale Definitions

Table 2. Scales that Measure the Risk and Protective Factors Shown in the Profiles

<i>School Domain Protective Factors</i>	
Opportunities for Prosocial Involvement	When young people are given more opportunities to participate meaningfully in important activities at school, they are less likely to engage in drug use and other problem behaviors.
Rewards for Prosocial Involvement	When young people are recognized and rewarded for their contributions at school, they are less likely to be involved in substance use and other problem behaviors.
<i>Peer-Individual Risk Factors</i>	
Early Initiation of Antisocial Behavior and Early Initiation of Drug Use	Early onset of drug use predicts misuse of drugs. The earlier the onset of any drug use, the greater the involvement in other drug use and the greater frequency of use. Onset of drug use prior to the age of 15 is a consistent predictor of drug abuse, and a later age of onset of drug use has been shown to predict lower drug involvement and a greater probability of discontinuation of use.
Attitudes Favorable Toward Antisocial Behavior and Attitudes Favorable Toward Drug Use	During the elementary school years, most children express anti-drug, anti-crim, and pro-social attitudes and have difficulty imagining why people use drugs or engage in antisocial behaviors. However, in middle school, as more youth are exposed to others who use drugs and engage in antisocial behavior, their attitudes often shift toward greater acceptance of these behaviors. Youth who express positive attitudes toward drug use and antisocial behavior are more likely to engage in a variety of problem behaviors, including drug use.
Perceived Risk of Drug Use	Young people who do not perceive drug use to be risky are far more likely to engage in drug use.
Interaction with Antisocial Peers	Young people who associate with peers who engage in problem behaviors are at higher risk for engaging in antisocial behavior themselves.
Friends' Use of Drugs	Young people who associate with peers who engage in alcohol or substance abuse are much more likely to engage in the same behavior. Peer drug use has consistently been found to be among the strongest predictors of substance use among youth. Even when young people come from well-managed families and do not experience other risk factors, spending time with friends who use drugs greatly increases the risk of that problem developing.
Rewards for Antisocial Behavior	Young people who receive rewards for their antisocial behavior are at higher risk for engaging further in antisocial behavior and substance use.
Depressive Symptoms	Young people who are depressed are overrepresented in the criminal justice system and are more likely to use drugs. Survey research and other studies have shown a link between depression and youth problem behaviors.
Gang Involvement	Youth who belong to gangs are more at risk for antisocial behavior and drug use.
<i>Peer-Individual Protective Factors</i>	
Religiosity	Young people who regularly attend religious services are less likely to engage in problem behaviors.
Belief in the Moral Order	Young people who have a belief in what is "right" or "wrong" are less likely to use drugs.
Interaction with Prosocial Peers	Young people who associate with peers who engage in prosocial behavior are more protected from engaging in antisocial behavior and substance use.
Prosocial Involvement	Participation in positive school and community activities helps provide protection for youth.
Rewards for Prosocial Involvement	Young people who are rewarded for working hard in school and the community are less likely to engage in problem behavior.

Data Tables

Table 3. Number of Students Who Completed the Survey

Number of Youth	Grade 7		Grade 8			Grade 9
	2015	2012	2014	2015	MTF 2014	2015
	963	1056	861	890	†	994

Table 4. Percentage of Students Who Used ATODs During Their Lifetime

Substance	Grade 7	Grade 8				Grade 9	
	2015	2012	2014	2015	MTF 2014	2015	
Alcohol	11.7	30.8	23.1	22.7	26.8	30.7	
Cigarettes	6.6	19.8	14.1	12.2	13.5	16.4	
Chewing Tobacco	3.6	9.3	5.6	4.4	8.0	8.3	
Marijuana	2.9	12.1	10.3	7.5	15.6	15.4	
Inhalants	4.3	9.8	7.0	5.6	10.8	4.7	
Hallucinogens	0.1	1.7	1.9	0.7	2.0	2.4	
Cocaine	0.6	1.7	1.0	0.6	1.8	1.0	
Methamphetamines	0.2	1.2	0.5	0.2	1.0	0.5	
Amphetamines	0.9	2.4	2.5	1.3	6.7	2.9	
Sedatives	2.6	4.3	2.6	3.5	5.5	4.3	
Tranquilizers	0.6	1.1	1.1	1.3	2.9	2.2	
Heroin	0.2	0.8	0.7	0.6	0.9	0.4	
Other Narcotics	0.5	2.2	1.9	0.6	2.3	2.3	
Ecstasy	0.2	1.0	0.7	0.4	1.4	1.3	
Synthetic Marijuana **	0.7	n/a	n/a	1.7	n/a	5.1	

† See the Monitoring The Future (MTF) website (www.monitoringthefuture.org).

MTF only surveys grades 8, 10 and 12.

** MTF either does not collect data on that substance at all, or not at that level.

Data Tables

Table 3. (continued) Number of Students Who Completed the Survey

Number of Youth	Grade 10				Grade 11	Grade 12				Total			
	2012	2014	2015	MTF 2014	2015	2012	2014	2015	MTF 2014	2012	2014	2015	MTF 2014
	964	724	920	†	860	715	518	944	†	2735	2103	5571	†

Table 4. (continued) Percentage of Students Who Used ATODs During Their Lifetime

Substance	Grade 10				Grade 11			Grade 12				Total			
	2012	2014	2015	MTF 2014	2012	2014	2015	2012	2014	2015	MTF 2014	2012	2014	2015	MTF 2014
Alcohol	60.4	49.7	43.8	49.3	51.8	72.2	61.1	59.0	66.0	52.2	41.5	36.4	46.4		
Cigarettes	29.4	25.2	24.1	22.6	24.8	39.3	26.4	27.8	34.4	28.4	20.9	18.6	22.9		
Chewing Tobacco	14.3	14.0	12.2	13.6	14.7	22.2	15.7	19.6	15.1	14.5	10.9	10.4	12.1		
Marijuana	32.6	25.7	25.8	33.7	30.7	43.6	35.4	33.8	44.4	27.6	21.6	19.2	30.5		
Inhalants	7.3	5.2	4.8	8.7	4.0	7.1	5.3	3.9	6.5	8.2	6.0	4.6	8.8		
Hallucinogens	4.4	4.2	3.4	5.0	5.3	11.3	4.8	6.1	6.3	5.2	3.4	3.0	4.3		
Cocaine	2.7	1.6	1.7	2.6	2.5	7.0	3.8	3.7	4.6	3.4	1.9	1.7	2.9		
Methamphetamines	1.4	0.7	0.9	1.4	1.0	2.5	1.1	0.9	1.9	1.6	0.7	0.6	1.4		
Amphetamines	9.1	4.4	4.9	10.6	3.9	13.1	6.3	6.7	12.1	7.5	4.1	3.5	9.7		
Sedatives	7.3	3.9	6.7	8.1	5.3	8.2	3.4	6.5	6.8	6.4	3.2	4.8	n/a		
Tranquilizers	4.0	2.0	4.5	5.8	4.6	7.3	4.0	6.2	7.4	3.7	2.1	3.2	5.3		
Heroin	1.1	0.7	0.8	0.9	0.8	1.9	0.6	0.8	1.0	1.2	0.7	0.6	0.9		
Other Narcotics	9.2	4.7	6.1	6.8	6.1	15.4	6.3	6.9	9.5	8.1	3.9	3.7	n/a		
Ecstasy	4.2	2.5	2.2	3.7	3.0	8.9	4.0	3.9	5.6	4.2	2.1	1.8	3.5		
Synthetic Marijuana **	n/a	n/a	7.8	n/a	6.6	n/a	n/a	6.3	n/a	n/a	n/a	4.7	n/a		

† See the Monitoring The Future (MTF) website (www.monitoringthefuture.org)
MTF only surveys grades 8, 10 and 12.
** MTF either does not collect data on that substance at all, or not at that level.

Data Tables

Table 5. Percentage of Students Who Used ATODs During The Past 30 Days

Substance	Grade 7		Grade 8					Grade 9	
	2015	2012	2014	2015	MTF 2014	2015	2015		
Alcohol	4.2	13.6	8.7	8.4	9.0	13.7			
Cigarettes	1.6	5.7	4.9	2.7	4.0	4.4			
Chewing Tobacco	1.2	3.7	2.0	2.6	3.0	3.5			
Marijuana	1.3	5.2	5.6	3.3	6.5	9.4			
Inhalants	1.5	3.9	3.1	2.2	2.2	1.7			
Hallucinogens	0.1	0.4	0.7	0.1	0.5	1.3			
Cocaine	0.2	0.5	0.2	0.0	0.5	0.6			
Methamphetamines	0.0	0.4	0.1	0.1	0.2	0.4			
Amphetamines	0.5	1.2	1.1	0.5	2.1	1.3			
Sedatives	0.5	1.8	1.2	1.6	1.6	1.9			
Tranquilizers	0.1	0.4	0.4	0.4	0.8	1.0			
Heroin	0.0	0.3	0.2	0.1	0.3	0.2			
Other Narcotics	0.2	0.8	0.5	0.1	0.7	0.9			
Ecstasy	0.1	0.3	0.4	0.1	0.4	0.4			
Synthetic Marijuana	0.5	n/a	n/a	1.0	4.4	2.8			
Energy Drinks **	18.6	n/a	n/a	26.1	n/a	28.8			
Caffeine Pills **	0.6	n/a	n/a	0.4	n/a	2.5			

Table 6. Percentage of Students With Problem ATOD Use

Problem Use	2016		2012		2014		2015		MTF 2014	2015
	2016	2012	2014	2015	2016	2012	2014	2015	2016	2015
Binge Drinking	2.3	6.4	5.4	3.7	4.1	7.0				
1/2 Pack of Cigarettes/Day	0.4	0.7	0.9	0.5	0.5	0.8				
Alcohol and Driving	2015	2012	2014	2015	2016	2015				
Drinking and Driving	0.6	1.1	1.0	0.8	5.6	0.8				
Riding with a Drinking Driver	12.6	15.2	10.0	11.7	22.3	14.2				

† See the Monitoring The Future (MTF) website (www.monitoringthefuture.org)
MTF only surveys grades 8, 10 and 12.
** MTF either does not collect data on that substance at all, or not at that level.

Data Tables

Table 5. (continued) Percentage of Students Who Used ATODs During The Past 30 Days

Substance	Grade 10				Grade 11		Grade 12				Total			
	2012	2014	2015	MTF 2014	2015	2012	2014	2015	MTF 2014	2012	2014	2015	MTF 2014	
Alcohol	33.8	22.7	24.5	23.5	25.4	48.5	36.7	34.9	37.4	29.9	20.3	18.5	22.6	
Cigarettes	11.9	8.7	9.0	7.2	8.3	17.1	11.1	11.0	13.6	10.9	7.7	6.2	8.0	
Chewing Tobacco	6.7	6.2	6.6	5.3	6.1	9.0	6.5	10.1	8.4	6.2	4.5	5.0	5.4	
Marijuana	18.2	13.1	15.9	16.6	16.1	25.7	16.2	19.1	21.2	15.1	10.7	10.8	14.4	
Inhalants	2.4	1.3	1.3	1.1	0.5	2.0	1.5	1.3	0.7	2.9	2.1	1.4	1.4	
Hallucinogens	0.7	0.4	1.2	1.2	1.6	3.3	1.3	2.5	1.5	1.3	0.8	1.1	1.0	
Cocaine	0.9	0.6	0.6	0.6	0.6	1.6	1.9	0.9	1.0	0.9	0.8	0.5	0.7	
Methamphetamines	0.4	0.3	0.2	0.3	0.3	0.6	0.4	0.1	0.5	0.5	0.3	0.2	0.3	
Amphetamines	3.7	1.8	1.6	3.7	1.3	5.0	1.9	2.4	3.8	3.1	1.5	1.3	3.2	
Sedatives	3.1	1.9	2.8	2.3	1.9	2.6	1.1	2.7	2.0	2.5	1.4	1.9	2.0	
Tranquilizers	1.0	0.1	1.6	1.6	2.0	2.2	0.8	2.1	2.1	1.1	0.4	1.2	1.5	
Heroin	0.4	0.3	0.0	0.4	0.1	0.6	0.6	0.5	0.4	0.4	0.4	0.2	0.3	
Other Narcotics	4.2	1.8	1.4	1.7	2.4	5.1	1.3	2.5	2.2	3.1	1.1	1.2	1.4	
Ecstasy	1.2	0.6	1.0	0.8	0.9	2.9	0.8	0.7	1.4	1.3	0.6	0.6	0.8	
Synthetic Marijuana	n/a	n/a	3.9	6.8	2.7	n/a	n/a	1.7	2.7	n/a	n/a	2.1	4.8	
Energy Drinks **	n/a	n/a	32.9	n/a	30.2	n/a	n/a	32.7	n/a	n/a	n/a	28.2	n/a	
Caffeine Pills **	n/a	n/a	2.0	n/a	4.0	n/a	n/a	4.6	n/a	n/a	n/a	2.3	n/a	

Table 6. (continued) Percentage of Students With Problem ATOD Use

Problem Use	2012	2014	2015	MTF 2014	2015	2012	2014	2015	MTF 2014	2012	2014	2015	MTF 2014
Binge Drinking	16.5	14.1	14.0	12.6	15.3	29.3	20.1	24.0	19.4	16.0	12.0	11.0	11.7
1/2 Pack of Cigarettes/Day	2.2	1.9	2.0	1.2	1.3	3.9	3.6	3.8	2.6	2.1	1.9	1.4	1.4
Alcohol and Driving	2012	2014	2015	BH Norm	2015	2012	2014	2015	BH Norm	2012	2014	2015	BH Norm
Drinking and Driving	1.9	1.4	1.6	5.3	3.1	5.8	5.1	3.9	11.8	2.6	2.1	1.8	7.4
Riding with a Drinking Driver	19.5	16.0	17.8	24.0	13.3	20.3	12.8	15.3	24.1	18.1	12.7	14.2	23.4

† See the Monitoring The Future (MTF) website (www.monitoringthefuture.org)

MTF only surveys grades 8, 10 and 12

** MTF either does not collect data on that substance at all, or not at that level.

Data Tables

ATOD Questions

Cigarettes	"Have you ever smoked cigarettes?" And "How frequently have you smoked cigarettes during the past 30 days?"
Chewing Tobacco	"Have you ever used smokeless tobacco (chew, snuff, plug, dipping tobacco, or chewing tobacco)? " and "How frequently have you used smokeless tobacco during the past 30 days?"
Or (if row many occasions (if any) have you:	
Alcohol	had alcoholic beverages (beer, wine or hard liquor) to drink in your lifetime (or during the past 30 days) – more than just a few sips?
Marijuana	used marijuana (grass, pot) or hashish (hash, hash oil) in your lifetime (or during the past 30 days)?
Inhalants	sniffed glue, breathed the contents of an aerosol spray can, or inhaled other gases or sprays, in order to get high in your lifetime (or during the past 30 days)?
Hallucinogens	used LSD (acid) or other hallucinogens (like PCP, mescaline, peyote, "shrooms" or psilocybin) in your lifetime (or during the past 30 days)?
Cocaine	used cocaine (like cocaine powder) or "crack" (cocaine in chunk or rock form) in your lifetime (or during the past 30 days)?
Methamphetamines	used methamphetamines (meth, speed, crank, crystal meth) in your lifetime (or during the past 30 days)?
Amphetamines	used prescription stimulants or amphetamines (such as Adderall, Ritalin, or Dexedrine) without a doctor telling you to take them, in your lifetime (or during the past 30 days)?
Sedatives	used prescription sedatives including barbiturates or sleeping pills (such as phenobarbital, Tuinal, Seconal, Ambien, Lunesta, or Sonata) without a doctor telling you to take them, in your lifetime (or during the past 30 days)?
Tranquilizers	used prescription tranquilizers (such as Librium, Valium, Xanax, Alivan, Soma, or Klonopin) without a doctor telling you to take them, in your lifetime (or during the past 30 days)?
Heroin	used heroin in your lifetime (or during the past 30 days)?
Other Narcotics	used narcotic prescription drugs (such as OxyContin, methadone, morphine, codeine, Demerol, Vicodin, Percocet) without a doctor telling you to take them, in your lifetime (or during the past 30 days)?
Ecstasy	used MDMA (X.E. or ecstasy) in your lifetime (or during the past 30 days)?
Synthetic Marijuana	used "synthetic marijuana" ("K2", "Spice") to get high in your lifetime (or during the past 30 days)?
Energy Drinks	drank energy drinks with caffeine (like Red Bull, Monster, Rockstar, or 5-Hour-Energy during the past 30 days)?
Caffeine Pills	used caffeine pills (No-Doz, Vivarin, Dexatrim) during the past 30 days?
Problem Use	
Binge Drinking	Think back over the last two weeks. How many times have you had five or more alcoholic drinks in a row? (A "drink" is a glass of wine, a bottle of beer, a wine cooler, a shot glass of liquor, or a mixed drink.)
1/2 Pack of Cigarettes/Day	How frequently have you smoked cigarettes during the past 30 days?
Alcohol and Driving	
Drinking and Driving	During the past 30 days, how many times did you DRIVE a car or other vehicle when you had been drinking alcohol?
Riding with a Drinking Driver	During the past 30 days, how many times did you RIDE in a car or other vehicle driven by someone who had been drinking alcohol?

Data Tables

Table 7. Percentage of Students With Antisocial Behavior in the Past Year

How many times in the past year (12 months) have you: (One or more times)	Grade 7			Grade 8			Grade 9		
	2015	BH Norm		2012	2014	2015	BH Norm	2015	BH Norm
Been Suspended from School	5.1	10.6		9.7	7.7	9.1	13.4	9.3	12.8
Been Drunk or High at School	2.2	4.0		5.9	5.7	3.7	7.8	8.2	11.5
Sold Illegal Drugs	1.1	1.2		2.0	3.1	1.2	3.1	3.7	5.2
Stolen or Tried to Steal a Motor Vehicle	0.9	1.5		0.7	0.8	1.2	2.2	1.4	2.8
Been Arrested	1.0	3.1		3.0	3.0	1.9	4.8	3.4	5.8
Attacked Someone with the Idea of Seriously Hurting Them	6.2	10.8		9.5	8.2	6.8	12.9	8.1	12.4
Carried a Handgun	4.1	5.0		3.4	2.4	3.7	5.4	4.5	5.0
Carried a Handgun to School	0.4	0.6		0.2	0.1	0.5	0.9	0.5	1.2

Table 7. (Contd.) Percentage of Students With Antisocial Behavior in the Past Year

How many times in the past year (12 months) have you: (One or more times)	Grade 10				Grade 11			Grade 12			Total			
	2012	2014	2015	BH Norm	2015	BH Norm	2012	2014	2015	BH Norm	2012	2014	2015	BH Norm
Been Suspended from School	9.6	8.2	6.9	11.2	7.3	13.8	9.2	5.8	7.2	8.5	9.6	7.4	7.5	11.1
Been Drunk or High at School	15.5	10.7	12.5	14.7	11.3	19.7	20.6	12.5	15.0	17.3	13.1	9.1	8.8	13.2
Sold Illegal Drugs	7.9	5.8	5.5	7.2	5.7	9.6	10.1	6.8	6.4	8.6	6.2	4.9	3.9	6.3
Stolen or Tried to Steal a Motor Vehicle	1.0	2.2	0.9	2.7	2.0	3.0	2.4	1.8	1.3	2.0	1.3	1.6	1.3	2.3
Been Arrested	4.5	3.9	4.0	6.0	3.2	7.8	7.1	3.0	4.6	5.8	4.6	3.3	3.0	5.6
Attacked Someone with the Idea of Seriously Hurting Them	9.5	9.3	7.4	11.8	6.7	12.9	7.7	5.6	6.9	9.6	9.1	7.9	7.0	11.5
Carried a Handgun	4.2	3.4	3.0	5.5	3.3	6.6	3.8	4.2	4.5	5.5	3.8	3.2	3.9	5.5
Carried a Handgun to School	0.4	1.1	0.3	1.2	1.0	1.5	1.1	1.6	1.3	1.2	0.5	0.8	0.7	1.1

Data Tables

Table 8. Percentage of Students Gambling in the Past Year

How many times in the past year (12 months) have you: (<i>'A few times' or more</i>)	Grade 7		Grade 8					Grade 9	
	2015	BH Norm	2012	2014	2015	BH Norm	2015	BH Norm	
Gambled in the Past Year	19.1	42.6	35.6	27.6	20.7	45.5	23.6	45.6	
Bet on Cards	3.8	15.8	13.4	6.3	7.2	25.0	5.9	21.0	
Gambled on the Internet	1.3	4.0	4.4	1.1	2.1	4.1	1.6	4.9	
Bet on Sports	7.0	20.6	16.3	12.1	10.4	23.4	9.9	22.2	
Played the Lottery	9.4	20.4	18.3	14.9	10.1	19.1	11.3	23.5	
Bet on Games of Skill	4.2	14.8	11.9	9.8	5.9	19.6	6.0	16.5	
Bet on Video Poker	1.7	3.3	3.7	2.0	2.2	4.2	1.5	4.8	
Bet on Dice	2.9	8.1	6.6	4.5	3.3	14.3	2.2	10.1	
Played Bingo for money	3.5	15.1	8.8	4.8	4.0	20.8	4.6	13.7	
Bet on Horses	2.5	7.4	3.7	2.3	2.5	5.8	2.1	8.6	
Gambled at a Casino	1.1	4.5	2.6	1.3	1.0	6.8	0.9	5.7	

Table 8. (Cont'd) Percentage of Students Gambling in the Past Year

How many times in the past year (12 months) have you: (A few times or more)	Grade 10				Grade 11		Grade 12				Total			
	2012	2014	2015	BH Norm	2015	BH Norm	2012	2014	2015	BH Norm	2012	2014	2015	BH Norm
Gambled in the Past Year	36.1	25.6	25.2	43.8	21.2	47.1	35.7	27.8	25.7	43.8	35.8	27.0	22.7	44.4
Bet on Cards	12.9	6.7	8.1	24.7	7.7	24.6	14.6	9.9	10.4	24.7	13.6	7.3	7.2	24.7
Gambled on the Internet	3.0	3.2	1.5	4.1	2.3	5.3	5.3	2.9	3.5	4.1	4.1	2.3	2.0	4.1
Bet on Sports	18.8	12.1	11.0	23.0	9.3	22.9	13.2	12.2	11.2	21.6	16.4	12.1	9.8	22.6
Played the Lottery	17.8	13.1	14.7	18.3	11.4	22.4	19.3	17.7	13.8	20.8	18.4	15.0	11.8	19.5
Bet on Games of Skill	11.3	7.4	6.3	18.4	5.3	17.4	8.9	6.8	7.4	16.9	10.9	8.3	5.9	18.2
Bet on Video Poker	3.9	3.4	2.3	4.0	1.8	4.9	5.3	4.7	3.0	4.5	4.2	3.1	2.1	4.3
Bet on Dice	4.3	4.5	3.3	12.5	2.3	11.5	5.6	3.7	4.3	11.1	5.5	4.3	3.0	12.5
Played Bingo for money	4.8	5.0	4.1	15.9	2.9	12.2	6.6	4.7	3.3	11.8	6.7	4.8	3.7	16.1
Bet on Horses	3.2	3.8	2.5	5.5	1.1	7.0	3.0	4.4	2.2	5.2	3.3	3.3	2.2	5.5
Gambled at a Casino	1.6	2.9	1.8	7.7	1.6	5.9	3.0	4.9	3.5	9.3	2.3	2.7	1.7	7.9

Data Tables

Table 9: Percentage of Students Reporting Protection

Protective Factors	Grade 7		Grade 8				Grade 9	
	2015	BH Norm	2012	2014	2015	BH Norm	2015	BH Norm
Community Domain								
Rewards for Prosocial Involvement	34.1	56.1	45.3	35.3	34.9	69.5	33.9	65.8
Family Domain								
Family Attachment	46.4	60.7	50.9	53.6	55.6	54.8	59.3	58.5
Opportunity for Prosocial Involvement	49.2	49.9	63.4	57.2	63.5	62.5	59.5	56.9
Rewards for Prosocial Involvement	56.3	55.5	43.3	46.8	48.4	61.9	56.2	55.5
School Domain								
Opportunity for Prosocial Involvement	72.1	66.0	61.7	62.4	64.4	62.1	66.3	64.7
Rewards for Prosocial Involvement	59.0	62.2	51.7	49.2	54.5	56.2	66.8	51.0
Peer-Individual Domain								
Religiosity	22.4	50.8	31.7	29.7	32.0	53.7	20.3	44.8
Belief in the Moral Order	58.1	53.6	70.8	75.8	76.1	52.1	68.0	59.5
Interaction with Prosocial Peers	50.2	60.1	44.9	50.0	43.8	54.9	51.1	54.9
Prosocial Involvement	53.5	63.1	58.3	59.0	57.0	58.1	63.4	65.0
Rewards for Prosocial Involvement	49.6	58.4	45.3	51.9	54.4	50.9	51.4	44.9
High Protection								
High Protection Youth **	66.1	64.5	59.8	59.7	61.2	44.9	61.1	54.0

** High Protection Youth are defined as the percentage of students who have more than a specified number of protective factors operating in their lives: (6th and 7th grades: 3 or more protective factors; 8th-12th grades: 4 or more factors)

Data Tables

Table 9. (Cont'd.) Percentage of Students Reporting Protection

Protective Factors	Grade 10				Grade 11		Grade 12			Total				
	2012	2014	2015	BH Norm	2015	BH Norm	2012	2014	2015	BH Norm	2012	2014	2015	BH Norm
Community Domain														
Rewards for Prosocial Involvement	37.7	31.2	30.5	65.2	29.3	63.6	39.9	32.3	33.0	64.6	41.0	33.2	32.6	66.4
Family Domain														
Family Attachment	55.4	55.4	59.2	57.1	56.5	55.2	57.2	57.1	56.6	57.9	54.2	55.0	55.7	56.6
Opportunity for Prosocial Involvement	55.1	55.7	54.1	56.2	56.1	53.8	55.6	56.1	54.8	56.2	58.4	56.4	56.2	58.1
Rewards for Prosocial Involvement	48.6	47.8	52.0	54.3	50.6	50.5	48.8	52.7	47.1	54.0	46.7	48.5	51.8	56.6
School Domain														
Opportunity for Prosocial Involvement	63.8	59.4	56.3	63.3	61.9	60.8	61.5	66.1	60.6	64.9	62.4	62.3	63.7	63.4
Rewards for Prosocial Involvement	69.1	61.0	61.7	48.0	60.2	48.5	65.8	71.7	66.5	50.3	61.6	58.8	61.6	51.5
Peer-Individual Domain														
Religiosity	30.5	24.6	28.1	48.4	23.1	39.6	24.9	26.7	23.4	42.9	29.5	27.2	24.8	48.3
Belief in the Moral Order	58.4	62.4	63.2	54.6	65.3	52.6	58.5	63.1	59.5	55.6	63.1	68.1	65.0	54.1
Interaction with Prosocial Peers	44.8	42.0	43.7	55.4	44.4	53.2	44.7	45.8	44.8	57.3	44.8	46.2	46.5	55.8
Prosocial Involvement	61.7	53.7	61.1	65.4	55.6	66.4	58.9	62.2	57.6	64.2	59.7	58.0	58.1	62.6
Rewards for Prosocial Involvement	51.5	52.6	60.5	48.8	45.0	48.0	42.0	45.9	49.3	52.2	46.6	50.7	51.7	50.6
High Protection														
High Protection Youth **	64.3	57.2	60.1	46.8	57.2	52.0	61.4	59.1	56.5	47.4	61.8	58.7	60.4	46.6

** High Protection Youth are defined as the percentage of students who have more than a specified number of protective factors operating in their lives: (6th and 7th grades: 3 or more protective factors; 8th-12th grades: 4 or more factors).

Data Tables

Table 10. Percentage of Students Reporting Risk

Risk Factors	Grade 7		Grade 8				Grade 9	
	2015	BH Norm	2012	2014	2015	BH Norm	2015	BH Norm
Community Domain								
Laws & Norms Favor Drug Use	37.9	38.4	39.9	37.1	37.5	40.0	43.9	42.7
Perceived Availability of Drugs	25.0	46.7	32.4	27.3	25.8	45.4	31.6	46.1
Family Domain								
Poor Family Management	42.4	45.4	41.6	38.1	39.4	40.4	35.8	46.5
Family Conflict	43.4	30.5	35.9	31.9	35.7	35.3	38.1	38.7
Sibling Drug Use	28.5	19.2	37.3	38.2	36.8	26.7	25.8	33.3
Exposure to Adult Antisocial Behavior	39.6	38.0	39.6	32.5	33.2	39.9	22.0	39.6
Parent Attitudes Favor Antisocial Behavior	31.1	41.8	45.2	37.6	42.2	49.1	40.0	52.2
Parent Attitudes Favor Drug Use	12.8	17.4	23.3	22.4	21.7	23.7	26.8	34.5
School Domain								
Academic Failure	36.3	40.0	41.7	40.0	38.6	41.1	36.4	39.5
Low Commitment to School	45.9	39.7	45.7	48.2	49.8	45.1	49.4	45.8
Peer-Individual Domain								
Early Initiation of Antisocial Behavior	16.6	26.9	27.3	24.9	22.8	32.2	25.6	32.1
Early Initiation of Drug Use	23.7	31.7	32.3	24.8	23.6	42.8	25.5	40.4
Attitudes Favorable to Antisocial Behavior	28.2	37.8	32.4	26.7	25.4	46.6	29.8	41.2
Attitudes Favorable to Drug Use	19.5	29.9	32.3	24.3	22.8	43.7	26.7	47.4
Perceived Risk of Drug Use	49.5	34.3	45.1	45.2	45.6	37.9	51.3	48.0
Interaction with Antisocial Peers	15.5	23.4	27.1	23.8	21.3	30.0	24.4	30.5
Friend's Use of Drugs	22.2	31.1	36.1	27.6	24.5	39.2	24.6	38.8
Rewards for Antisocial Behavior	30.1	32.3	37.9	32.2	28.2	43.2	42.3	49.1
Depressive Symptoms	34.2	32.6	36.2	37.6	36.7	34.8	41.0	36.9
Gang Involvement	5.4	6.7	6.5	8.0	5.4	6.9	6.2	7.3
High Risk								
High Risk Youth *	29.5	35.9	41.5	37.6	36.3	40.3	37.8	49.5

* High Risk Youth are defined as the percentage of students who have more than a specified number of risk factors operating in their lives (6th grade: 5 or more risk factors; 7th-9th grades: 6 or more factors; 10th-12th grades: 7 or more factors)

Data Tables

Table 10. (Contd.) Percentage of Students Reporting Risk

Risk Factors	Grade 10				Grade 11		Grade 12				Total			
	2012	2014	2015	BH Norm	2015	BH Norm	2012	2014	2015	BH Norm	2012	2014	2015	BH Norm
Community Domain														
Laws & Norms Favor Drug Use	45.9	42.7	45.0	48.4	52.2	49.9	59.5	58.0	56.7	48.1	47.5	44.1	45.7	45.5
Perceived Availability of Drugs	37.1	27.5	30.7	47.4	25.5	46.1	40.8	28.8	27.7	41.0	36.4	27.7	27.8	44.7
Family Domain														
Poor Family Management	37.7	36.2	32.1	40.0	35.7	48.2	40.1	37.7	38.1	41.2	39.8	37.4	37.2	40.7
Family Conflict	43.2	41.5	36.8	39.9	43.5	41.2	39.5	37.6	39.7	38.0	39.5	36.5	39.5	37.8
Sibling Drug Use	41.3	31.1	33.2	39.4	36.4	45.7	43.3	45.4	39.0	47.4	40.4	37.5	33.1	37.7
Exposure to Adult Antisocial Behavior	49.6	43.5	38.2	46.6	33.6	43.2	48.3	34.7	33.2	42.9	45.7	36.8	33.1	43.1
Parent Attitudes Favor Antisocial Behavior	55.1	45.8	46.7	53.5	44.0	55.4	55.0	44.3	46.4	52.9	51.4	42.0	41.8	51.8
Parent Attitudes Favor Drug Use	48.2	40.2	40.1	39.6	32.5	50.8	48.2	40.6	37.9	40.3	39.0	32.9	28.8	34.3
School Domain														
Academic Failure	38.7	44.4	42.3	39.8	42.0	39.4	39.1	35.6	42.1	37.9	39.9	40.4	39.5	39.7
Low Commitment to School	40.5	45.1	50.3	41.1	49.0	46.2	43.2	41.6	48.3	43.6	43.2	45.5	48.8	43.3
Peer/Individual Domain														
Early Initiation of Antisocial Behavior	30.2	29.0	26.2	34.2	27.0	35.0	32.3	24.3	27.7	34.2	29.6	26.2	24.3	33.5
Early Initiation of Drug Use	35.3	28.4	27.2	41.4	27.5	42.7	41.3	34.3	34.5	41.8	35.7	28.4	27.0	41.9
Attitudes Favorable to Antisocial Behavior	41.4	35.6	31.4	40.8	30.9	43.5	41.6	31.4	34.1	39.0	38.0	30.9	30.0	42.1
Attitudes Favorable to Drug Use	48.8	39.7	35.0	45.3	33.8	43.8	50.9	39.5	40.0	46.9	43.0	33.3	29.6	45.1
Perceived Risk of Drug Use	64.6	63.2	61.4	51.9	54.5	47.6	60.5	58.8	61.0	47.4	56.1	54.7	53.9	45.5
Interaction with Antisocial Peers	31.5	25.7	22.1	31.3	23.2	36.8	29.0	18.0	23.7	29.6	29.1	23.0	21.7	30.3
Friends Use of Drugs	40.1	31.2	30.4	40.4	31.4	43.9	36.9	27.2	29.5	44.7	37.7	28.7	27.0	41.3
Rewards for Antisocial Behavior	57.2	44.9	43.7	46.7	48.7	50.2	62.8	48.8	48.0	51.5	51.3	40.7	40.1	46.9
Depressive Symptoms	39.1	41.5	41.5	37.8	40.6	37.8	37.3	32.6	39.5	33.4	37.5	37.7	38.9	35.4
Gang Involvement	6.4	4.3	4.2	5.9	5.5	7.5	5.0	5.1	5.3	5.2	6.1	6.0	5.3	6.0
High Risk														
High Risk Youth *	49.5	39.9	39.6	42.0	37.9	48.6	51.2	39.0	38.7	41.9	46.8	38.8	36.6	41.5

* High Risk Youth are defined as the percentage of students who have more than a specified number of risk factors operating in their lives (6th grade: 5 or more risk factors; 7th-9th grades: 6 or more factors; 10th-12th grades: 7 or more factors)

Student Alcohol Table

Table 11. Sources and Places of Student Alcohol Use

If you drank alcohol (not just a sip or taste) in the past year (12 months), how did you get it?	Grade 7		Grade 8			Grade 9		Grade 10			Grade 11		Grade 12			Total	
	2015	2012	2014	2015	2015	2012	2014	2015	2015	2015	2012	2014	2015	2012	2014	2015	2015
Sample size *	104	258	149	143	244	520	300	357	380	489	275	457	1,247	724	1,702		
I bought it myself from a store.	4.0	1.6	2.0	2.1	2.0	2.5	4.7	1.4	5.8	5.1	6.2	6.6	3.3	4.7	4.1		
I got it at a party.	30.7	43.4	48.3	37.1	50.4	65.4	58.3	54.8	64.7	76.8	72.4	69.8	65.1	61.6	57.6		
I gave someone else money to buy it for me.	12.9	15.5	12.8	6.3	15.6	33.8	24.3	28.3	36.8	56.1	47.3	43.9	38.4	30.7	29.9		
I got it from someone I know age 21 or older.	35.6	46.9	49.7	37.8	47.5	54.6	49.0	45.2	59.7	73.8	64.4	64.2	60.2	55.0	52.8		
I got it from someone I know under age 21.	25.7	27.5	22.8	27.3	31.6	44.4	41.3	36.2	36.8	43.5	38.2	40.0	40.6	36.3	35.4		
I got it from a family member or relative other than my parents.	25.7	39.9	37.6	37.1	38.1	39.6	39.0	37.9	37.9	40.1	41.8	40.9	39.9	39.8	38.0		
I got it from home with my parents' permission.	48.5	38.0	37.6	48.3	38.5	38.3	45.0	39.2	43.2	44.1	49.1	43.7	40.4	45.0	42.5		
I got it from home without my parents' permission.	25.7	43.8	36.9	36.4	34.8	45.8	44.3	40.1	35.5	34.3	34.5	36.2	41.1	39.1	36.1		
I got it at work.	4.0	n/a	5.4	0.7	1.6	n/a	4.3	1.6	4.7	n/a	2.9	4.3	n/a	4.0	3.1		
I bought it over the internet.	5.0	n/a	1.3	0.7	1.6	n/a	3.0	0.8	1.8	n/a	1.8	2.1	n/a	2.2	1.8		
I got it in another way.	24.8	22.5	26.8	16.1	12.3	16.7	13.0	14.4	8.2	13.4	6.9	8.4	16.7	13.5	11.8		
During the past year (12 months) did you drink alcohol at any of the following places?	Grade 7		Grade 8			Grade 9		Grade 10			Grade 11		Grade 12			Total	
	2015	2012	2014	2015	2015	2012	2014	2015	2015	2015	2012	2014	2015	2012	2014	2015	
Sample size *	112	261	159	163	252	622	290	360	359	480	275	458	1,248	724	1,899		
At my home or someone else's home without any parent permission.	27.7	50.2	44.7	47.2	52.4	62.5	53.1	60.0	57.7	58.5	56.0	56.7	58.4	52.3	54.1		
At my home with my parent's permission.	64.3	50.2	48.4	60.1	49.6	47.1	53.8	47.8	50.1	54.1	56.7	54.5	50.4	53.7	52.6		
At someone else's home with their parent's permission.	14.3	19.5	24.5	16.0	23.0	33.7	35.5	29.2	37.0	56.3	47.3	50.8	39.1	37.6	33.4		
At an open area like a park, beach, or back road.	17.0	24.1	23.9	9.2	21.4	28.7	28.3	27.8	28.7	43.0	27.3	30.2	33.1	26.9	25.2		
At public events such as a sporting event, festival, or concert.	11.6	13.8	9.4	6.7	15.1	17.4	16.2	17.8	15.3	23.0	18.5	23.4	18.7	15.6	16.9		
At a restaurant, bar, or a nightclub.	4.5	8.4	6.9	3.1	6.7	8.0	9.0	7.5	7.8	13.0	10.2	13.9	10.0	9.0	8.5		
In a car.	8.9	14.2	8.8	6.1	11.5	20.1	21.4	18.9	19.5	28.5	24.7	27.8	22.0	19.9	18.4		
At a school dance, a game, or other event.	9.8	10.7	6.9	4.9	10.3	14.4	11.0	12.8	9.2	13.3	13.8	15.7	13.2	11.2	11.5		
At school during the day.	6.3	5.0	6.3	1.8	4.8	9.0	11.0	5.8	5.3	9.6	5.8	7.5	8.4	8.0	5.7		
Near school.	8.9	7.3	8.2	2.5	7.1	10.3	7.6	8.3	6.4	10.2	8.7	9.5	9.7	8.1	7.5		
In another place.	19.6	27.2	27.0	16.0	23.0	29.3	18.3	21.9	22.0	27.4	20.4	23.2	28.2	21.0	21.7		

* Sample size represents the number of youth who answered the question, not including students reporting no use in the past year. In the case of smaller sample sizes, caution should be exercised before generalizing results and yearly trends to the entire community.

CSAP Questions

Table 12. CSAP Questions

Question	Response	Grade 7		Grade 8		Grade 9		Grade 10		Grade 11		Grade 12		Total	
		Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent
For the following for questions, during the past 12 months, have you talked with at least one of your parents (by parents, we mean either your biological parents, adoptive parents, stepparents, foster parents, or other adult caregivers whether or not they live with you) about:															
The dangers of underage drinking?	Yes	424	58.3	417	55.0	449	53.2	433	52.5	375	49.5	396	50.1	2494	53.0
	No	303	41.7	341	45.0	395	46.8	392	47.5	383	50.5	395	49.9	2209	47.0
The dangers of tobacco use?	Yes	452	62.0	431	56.9	457	54.1	456	55.4	392	51.8	405	51.2	2693	55.1
	No	277	38.0	326	43.1	388	45.9	367	44.6	365	48.2	386	48.8	2109	44.9
When parents find out their kids have been drinking they may discuss it, take away privileges, add chores, take away cell phones, use of the car, etc. In the past 12 months, if your parents found out you were drinking, how did they usually respond? (Select one option).	I didn't drink.	657	87.6	639	82.5	664	78.7	536	65.4	477	62.6	462	57.2	3435	72.2
	I drank but I was not caught.	15	2.0	51	6.6	72	8.5	113	13.8	120	15.7	124	15.3	495	10.4
	I was caught but there were no consequences.	8	1.1	18	2.3	29	3.4	47	5.7	72	9.4	114	14.1	288	6.1
	There were minor consequences.	9	1.2	11	1.4	20	2.4	45	5.5	48	6.3	59	7.3	192	4.0
	There were major consequences.	61	8.1	56	7.2	59	7.0	78	9.5	45	5.9	49	6.1	348	7.3
If the police caught you drinking, which of the following would most likely happen? (Select one option).	There would be no consequence.	44	6.1	44	5.7	56	6.7	66	8.1	62	8.2	72	9.0	344	7.3
	I would be given a warning and then let go.	48	6.6	54	7.0	90	10.7	91	11.2	89	11.8	120	14.9	492	10.5
	I would be taken home to my parents.	366	50.5	426	55.6	447	53.1	422	52.0	382	50.5	390	48.5	2433	51.7
	I would be arrested but would get no penalty.	100	13.8	85	11.1	95	11.3	87	10.7	80	10.6	66	8.2	513	10.9
	I would be arrested and the court would impose a penalty.	167	23.0	157	20.5	154	18.3	146	18.0	143	18.9	156	19.4	923	19.6
For the following for questions, during the past 12 months, how many times has each of the following things happened?															
You had problems at school or work because you had been drinking.	Never	769	99.6	767	98.5	849	98.6	812	97.8	751	98.2	790	97.1	4738	98.3
	Once	2	0.3	7	0.9	7	0.8	7	0.8	7	0.9	17	2.1	47	1.0
	Twice	0	0.0	1	0.1	2	0.2	5	0.6	2	0.3	4	0.5	14	0.3
	Three or four times	1	0.1	0	0.0	0	0.0	1	0.1	2	0.3	2	0.2	6	0.1
	Five or more times	0	0.0	4	0.5	3	0.3	5	0.6	3	0.4	1	0.1	16	0.3

CSAP Questions

Table 12. (Contd.) CSAP Questions

Question	Response	Grade 7		Grade 8		Grade 9		Grade 10		Grade 11		Grade 12		Total	
		Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent
For the following for questions, during the past 12 months, how many times has each of the following things happened?															
You had problems with your friends because you had been drinking.	Never	769	99.7	764	98.2	833	96.9	791	95.3	731	95.8	776	95.3	4664	96.8
	Once	0	0.0	10	1.3	19	2.2	25	3.0	21	2.8	21	2.6	96	2.0
	Twice	1	0.1	1	0.1	5	0.6	6	0.7	2	0.3	9	1.1	24	0.5
	Three or four times	0	0.0	0	0.0	2	0.2	2	0.2	5	0.7	4	0.5	13	0.3
	Five or more times	1	0.1	3	0.4	1	0.1	6	0.7	4	0.5	4	0.5	19	0.4
You had problems with someone you were dating because you had been drinking.	Never	768	99.9	765	98.3	827	96.6	779	94.0	734	96.2	759	93.5	4632	96.4
	Once	0	0.0	7	0.9	15	1.8	29	3.5	11	1.4	27	3.3	89	1.9
	Twice	0	0.0	2	0.3	10	1.2	9	1.1	12	1.6	9	1.1	42	0.9
	Three or four times	1	0.1	1	0.1	1	0.1	6	0.7	3	0.4	10	1.2	22	0.5
	Five or more times	0	0.0	3	0.4	3	0.4	6	0.7	3	0.4	7	0.9	22	0.5
You were hung over.	Never	761	99.3	748	96.0	797	92.9	693	83.7	631	82.7	646	79.6	4276	89.0
	Once	3	0.4	18	2.3	34	4.0	44	5.3	47	6.2	37	4.6	183	3.8
	Twice	1	0.1	2	0.3	15	1.7	29	3.5	23	3.0	51	6.3	121	2.5
	Three or four times	0	0.0	6	0.8	6	0.7	29	3.5	29	3.8	35	4.3	105	2.2
	Five or more times	1	0.1	5	0.6	6	0.7	33	4.0	33	4.3	43	5.3	121	2.5
You were sick to your stomach or threw up after drinking.	Never	759	99.6	754	97.0	820	95.8	746	90.0	669	87.7	693	85.6	4441	92.6
	Once	2	0.3	13	1.7	23	2.7	38	4.6	48	6.3	51	6.3	175	3.6
	Twice	0	0.0	4	0.5	7	0.8	16	1.9	22	2.9	32	4.0	81	1.7
	Three or four times	1	0.1	1	0.1	3	0.4	15	1.8	11	1.4	20	2.5	51	1.1
	Five or more times	0	0.0	5	0.6	3	0.4	14	1.7	13	1.7	14	1.7	49	1.0
You got into a sexual situation that you later regretted because you had been drinking.	Never	765	99.7	765	98.5	835	97.3	783	94.5	715	94.0	741	91.3	4604	95.8
	Once	1	0.1	6	0.8	17	2.0	27	3.3	24	3.2	41	5.0	116	2.4
	Twice	0	0.0	2	0.3	3	0.3	8	1.0	11	1.4	15	1.8	39	0.8
	Three or four times	0	0.0	0	0.0	1	0.1	2	0.2	2	0.3	10	1.2	15	0.3
	Five or more times	1	0.1	4	0.5	2	0.2	9	1.1	9	1.2	5	0.6	30	0.6
You got into a physical fight because you had been drinking.	Never	760	99.6	766	98.6	841	98.0	800	96.6	735	96.6	777	95.8	4679	97.5
	Once	2	0.3	6	0.8	8	0.9	18	2.2	13	1.7	15	1.8	62	1.3
	Twice	1	0.1	1	0.1	6	0.7	3	0.4	8	1.1	10	1.2	29	0.6
	Three or four times	0	0.0	0	0.0	2	0.2	2	0.2	3	0.4	4	0.5	11	0.2
	Five or more times	0	0.0	4	0.5	1	0.1	5	0.6	2	0.3	5	0.6	17	0.4
You were drunk at school or work.	Never	761	99.6	769	99.1	847	98.9	804	97.2	733	96.4	773	95.7	4687	97.8
	Once	2	0.3	2	0.3	7	0.8	13	1.6	10	1.3	20	2.5	54	1.1
	Twice	0	0.0	1	0.1	1	0.1	2	0.2	6	0.8	10	1.2	20	0.4
	Three or four times	0	0.0	0	0.0	0	0.0	1	0.1	3	0.4	3	0.4	7	0.1
	Five or more times	1	0.1	4	0.5	1	0.1	7	0.8	8	1.1	2	0.2	23	0.5

DFC and Youth Perception Tables

Table 13. Drug Free Communities Report *

Outcomes	Definition	Grade 7		Grade 8		Grade 9		Grade 10		Grade 11		Grade 12		Total †		Male		Female	
		Per.	Num.	Per.	Num.	Per.	Num.	Per.	Num.	Per.	Num.	Per.	Num.	Per.	Num.	Per.	Num.	Per.	Num.
How much do you think people risk harming themselves (physically or in other ways) if they: (Moderate risk or Great Risk)	take one or two drinks of an alcoholic beverage (beer, wine, liquor) nearly every day?	69.5	869	70.0	833	71.1	914	69.9	866	66.1	793	62.3	853	68.2	5128	65.5	2502	70.6	2597
	have five or more drinks of an alcoholic beverage once or twice a week?	76.6	864	77.4	831	79.0	916	73.3	870	70.1	797	68.6	850	74.3	5128	72.0	2503	76.4	2596
	smoke 1 or more packs of cigarettes per day.	82.0	867	79.0	835	82.2	917	81.2	867	82.5	798	79.8	856	81.1	5140	81.4	2506	80.8	2605
	smoke marijuana once or twice a week?	72.9	864	66.9	829	63.2	917	52.0	868	42.5	793	40.3	852	56.5	5123	53.4	2502	59.6	2592
	use prescription pain relievers that are not prescribed to them?	77.7	860	78.6	829	82.3	915	77.9	866	77.8	794	75.9	855	78.4	5119	77.8	2499	79.1	2591
How wrong do your parents feel it would be for YOU to: (Wrong or Very Wrong)	have one or two drinks of an alcoholic beverage nearly every day?	98.7	784	98.0	782	97.8	868	95.4	835	94.8	766	93.3	820	96.3	4855	95.5	2333	97.0	2493
	smoke cigarettes	98.7	788	97.2	783	96.2	867	95.7	839	94.3	769	92.1	818	95.7	4864	95.7	2341	95.6	2494
	smoke marijuana	98.2	786	95.8	784	93.7	867	90.3	838	85.8	768	86.6	818	91.7	4861	91.0	2338	92.5	2494
	use prescription pain relievers that are not prescribed to them?	97.8	787	97.6	782	96.7	869	95.7	838	95.8	769	94.9	819	96.4	4864	96.4	2339	96.4	2496
How wrong do your friends feel it would be for you to: (Wrong or Very Wrong)	have one or two drinks of an alcoholic beverage nearly every day?	92.7	924	90.2	855	84.8	954	74.4	887	70.8	815	66.3	898	80.0	5333	77.9	2612	82.1	2693
	smoke cigarettes	95.8	922	92.3	862	88.7	951	82.7	886	79.4	817	72.9	897	85.4	5325	84.2	2614	86.7	2683
	smoke marijuana	93.6	911	87.2	839	78.3	944	65.9	880	63.7	811	56.1	888	74.3	5273	72.5	2591	76.2	2655
	use prescription pain relievers that are not prescribed to them?	93.8	922	94.4	851	89.8	949	85.2	888	84.5	814	81.6	899	88.3	5323	88.1	2613	88.4	2682
How do you feel about someone your age having one or two drinks of an alcoholic beverage nearly every day?	Neither Approve nor Disapprove	16.1	140	21.0	172	21.4	197	24.0	207	31.8	253	31.8	271	24.2	1240	26.8	670	21.6	560
	Somewhat Disapprove	5.2	45	11.1	91	13.7	126	18.0	155	16.8	134	18.7	159	13.9	710	14.4	360	13.4	347
	Strongly Disapprove	65.7	572	58.1	475	53.3	491	47.1	406	42.0	334	39.2	334	51.0	2612	47.6	1187	54.6	1416
	Don't know or can't say	13.1	114	9.8	80	11.6	107	10.9	94	9.4	75	10.3	88	10.9	558	11.2	279	10.5	272
Past 30 day use of (at least one use in the Past 30 Days):	Alcohol	4.2	854	8.4	824	13.7	912	24.5	866	25.4	792	34.9	856	18.5	5104	17.4	2476	19.5	2600
	Cigarettes	1.6	849	2.7	814	4.4	901	9.0	866	8.3	784	11.0	840	6.2	5054	6.0	2448	6.3	2577
	Marijuana	1.3	851	3.3	823	9.4	909	15.9	866	16.1	785	19.1	850	10.8	5084	11.9	2463	9.7	2592
	Any Prescription Drug	1.3	852	2.1	817	3.3	908	4.3	867	4.9	788	5.8	848	3.6	5080	3.2	2457	4.0	2594
Average Age of Onset **	Alcohol	Age	Num.	Age	Num.	Age	Num.	Age	Num.	Age	Num.	Age	Num.	Age	Num.	Age	Num.	Age	Num.
	Cigarettes	11.0	191	11.6	281	12.4	352	13.0	463	13.7	475	14.0	581	13.0	2323	12.7	1126	13.2	1183
	Marijuana	10.8	80	11.4	111	12.0	173	12.8	230	13.1	231	13.6	288	12.6	1113	12.5	542	13.8	565
	Prescription Pain Relievers	11.3	29	12.3	69	12.9	151	13.4	232	14.0	283	14.3	342	13.7	1106	13.5	577	13.9	521
		10.7	13	11.8	12	12.7	18	13.2	38	14.5	31	15.1	36	13.6	148	13.9	71	13.3	76

*The "Num." column represents the sample size (the number of youth who answered the question). The "Per." column represents the percentage of youth in the sample answering the question as specified.

**For Average Age of Onset, "Num." represents the number of youth who reported any age of first use for the specified substance other than "Never Used."

†The "Total" column represents responses from students in all grades surveyed.

DFC and Youth Perception Tables

Table 14. Youth Perceptions of Substance Use

Now think about all the students in your grade at school. How many of them do you think:	Substance	Grade 7		Grade 8		Grade 9		Grade 10		Grade 11		Grade 12		Total	
		Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent	Number	Percent
a. smoke one or more cigarettes a day?	None (0%)	472	50.8	319	36.7	242	24.9	164	18.1	139	16.7	155	16.9	1491	27.5
	Few (1-10%)	250	26.9	297	34.1	234	24.0	243	26.9	181	21.7	192	20.9	1397	25.7
	Some (11-30%)	118	12.7	142	16.3	232	23.8	201	22.2	206	24.7	237	25.8	1136	20.9
	Half or less (31-50%)	49	5.3	58	6.7	117	12.0	150	16.6	136	16.3	167	18.2	677	12.5
	Half or more (51-70%)	27	2.9	35	4.0	94	9.7	91	10.1	114	13.7	100	10.9	461	8.5
	Most (71-90%)	5	0.5	12	1.4	40	4.1	41	4.5	44	5.3	53	5.8	195	3.6
b. drank alcohol sometime in the past month?	Almost All (91-100%)	9	1.0	7	0.8	14	1.4	14	1.5	14	1.7	13	1.4	71	1.3
	None (0%)	471	50.9	330	38.1	219	22.6	130	14.4	113	13.6	109	11.9	1372	25.3
	Few (1-10%)	247	26.7	261	30.1	205	21.1	118	13.0	70	8.4	72	7.9	973	18.0
	Some (11-30%)	115	12.4	147	17.0	192	19.8	167	18.5	143	17.2	138	15.1	902	16.7
	Half or less (31-50%)	52	5.6	68	7.9	151	15.6	179	19.8	167	20.0	150	16.4	767	14.2
	Half or more (51-70%)	24	2.6	32	3.7	122	12.6	156	17.2	162	19.4	175	19.1	671	12.4
c. used marijuana sometime in the past month?	Most (71-90%)	7	0.8	24	2.8	62	6.4	114	12.6	138	16.6	197	21.6	542	10.0
	Almost All (91-100%)	10	1.1	4	0.5	19	2.0	41	4.5	40	4.8	73	8.0	187	3.5
	None (0%)	570	61.6	380	43.9	246	25.4	159	17.6	122	14.6	136	14.8	1613	29.8
	Few (1-10%)	198	21.4	238	27.5	207	21.4	150	16.6	110	13.2	98	10.7	1001	18.5
	Some (11-30%)	61	6.6	104	12.0	158	16.3	162	18.0	149	17.9	165	18.0	799	14.8
	Half or less (31-50%)	37	4.0	61	7.1	129	13.3	144	16.0	151	18.1	153	16.7	675	12.5
d. used an illegal drug in the past month (not including marijuana)?	Half or more (51-70%)	29	3.1	31	3.6	90	9.3	126	14.0	115	13.8	153	16.7	544	10.1
	Most (71-90%)	18	1.9	32	3.7	98	10.1	94	10.4	119	14.3	153	16.7	514	9.5
	Almost All (91-100%)	13	1.4	19	2.2	40	4.1	67	7.4	68	8.2	58	6.3	265	4.9
	None (0%)	602	64.9	489	56.7	340	35.1	243	26.9	206	24.7	223	24.4	2103	38.9
	Few (1-10%)	190	20.5	216	25.0	310	32.0	303	33.6	266	31.9	291	31.8	1576	29.1
	Some (11-30%)	80	8.6	81	9.4	146	15.1	160	17.7	143	17.1	160	17.5	770	14.2
	Half or less (31-50%)	29	3.1	46	5.3	64	6.6	85	9.4	100	12.0	100	10.9	424	7.8
	Half or more (51-70%)	10	1.1	15	1.7	61	6.3	52	5.8	66	7.9	74	8.1	278	5.1
	Most (71-90%)	6	0.6	9	1.0	33	3.4	39	4.3	38	4.6	45	4.9	170	3.1
	Almost All (91-100%)	10	1.1	7	0.8	16	1.6	20	2.2	16	1.9	22	2.4	91	1.7

NATIONAL RESOURCES

Human Services (USDHHS)
Substance Abuse and Mental Health
Service Administration (SAMHSA)
1 Choke Cherry Rd., Rm. 8-1054
Rockville, Maryland 20857
240-276-2000
info@samhsa.hhs.org

(From this web-site, the programs and
services provided by the Center for
Substance Abuse Treatment, and Center for
Mental Health Services can be accessed)

Center for Substance Abuse Prevention

(CSAP)
1 Choke Cherry Rd., Ste 4-1057
Rockville, Maryland 20857
240-276-2420

info@samhsa.hhs.org

<http://prevention.samhsa.gov/>

CSAP's Centers for the Advancement of
Prevention Technologies (all five CSAP
Centers can be accessed through this web
site)

<http://capitus.samhsa.gov/home.cfm>

National Institutes of Health (NIH)

6001 Executive Blvd., Rm. 5213

Bethesda, Maryland 20892-9561

301-443-1124

<http://www.nida.nih.gov/>

STATE RESOURCES

New York State
Office of Alcoholism and Substance Abuse
Services (OASAS),
Division of Prevention and Treatment
prevention@oasas.state.ny.us
www.oasas.state.ny.us

This Report Was Prepared
by Bach Harrison L.L.C.
R. Steven Harrison, Ph.D.
R. Paris Bach-Harrison, B.F.A.
Taylor C. Bryant, B.A.
801-359-2064
<http://www.bach-harrison.com>